



Some Common Fixed Point Theorems in Polish Space Using New Type of Contractive Conditions

Rajesh Shrivastava¹, Rajendra Kumar Dubey² and Pankaj Tiwari¹

¹Department of Mathematics, Govt. Science and Comm. College Benjeer, Bhopal, M.P., INDIA

²Department of Mathematics, Govt. Science P.G. College, Rewa M.P., INDIA

Available online at: www.isca.in, www.isca.me

Received 12th October 2013, revised 2th November 2013, accepted 9th November 2013

Abstract

In this paper, we established Some Common Fixed Point Theorems for Random Operator in polish spaces, by using some new type of contractive condition. Our result is generalization of various known results.

Keywords: Polish space, random operator, random multivalued operator, random fixed point, measurable mapping.

AMS Subject Classification: 47H10, 54H25.

Introduction

Probabilistic functional analysis has emerged as one of the important mathematical disciplines in view of its role in analyzing Probabilistic models in the applied sciences. The study of fixed point of random operator forms a central topic in this area. Random fixed point theorem for contraction mappings in Polish spaces and random fixed point theorems are of fundamental importance in probabilistic functional analysis. There study was initiated by the Prague school of Probabilistics, in 1950, with their work of Spacek¹ and Hans^{2,3}. For example surveys are referring to Bharucha-Reid⁴ Itoh⁵ proved several random fixed point theorems and gave their applications to Random differential equations in Banach spaces. Random coincidence point theorems and random fixed point theorems are stochastic generalization of classical coincidence point theorems and classical fixed point theorems.

Random fixed point theorems are stochastic generalization of classical fixed point theorems. Itoh⁵ extended several well known fixed point theorems, thereafter, various stochastic aspects of Schauder's fixed point theorem have been studied by Sehgal and Singh⁶, Papageorgiou⁷, Lin⁸ and many authors. In a separable metric space, random fixed point theorems for contractive mappings were proved by Spacek¹, Hans^{2,3}. Afterwards, Beg and Shahzad⁹, Badshah and Sayyad studied the structure of common random fixed points and random coincidence points of a pair of compatible random operators and proved the random fixed point theorems for contraction random operators in Polish spaces.

Preliminaries: In this section, we give some definitions which are useful to prove our results.

Definition: 2.1: A metric space (X, d) is said to be a Polish Space, if it satisfying following conditions: i. X is complete, ii. X is separable,

Before we describe our next hierarchy of set of reals of ever increasing complexity, we would like to consider a class of metric spaces under which we can unify $2^\omega, \omega^\omega, \mathcal{R}$ and there products. This will be helpful in formulating this hierarchy. (as well as future ones) Recall that a metric space (X, d) is complete if whenever $(x_n: n \in \omega)$ is a sequence of member of X , such that for every $\epsilon > 0$ there is an N , such that $m, n \geq N$ implies $d(x_n, x_m) < \epsilon$, there is a single x in X such that $\lim_{n < \omega} x_n = x$. It is easy to see that $2^\omega, \omega^\omega$ are polish space, So in fact is ω under the discrete topology, whose metric is given by letting $d(x, y) = 1$ when $x \neq y$ and $d(x, y) = 0$ when $x = y$. Let (X, d) be a Polish space that is a separable complete metric space and (Ω, \mathcal{q}) be Measurable space. Let 2^X be a family of all subsets of X and $CB(X)$ denote the family of all nonempty bounded closed subsets of X . A mapping $T: \Omega \rightarrow 2^X$ is called measurable if for any open subset C of X , $T^{-1}(C) = \{\omega \in \Omega: f(\omega) \cap C \neq \emptyset\} \in \mathcal{q}$. A mapping $\xi: \Omega \rightarrow X$ is said to be measurable selector of a measurable mapping $T: \Omega \rightarrow 2^X$, if ξ is measurable and for any $\omega \in \Omega, \xi(\omega) \in T(\omega)$. A mapping $f: \Omega \times X \rightarrow X$ is called random operator, if for any $x \in X, f(\cdot, x)$ is measurable. A Mapping $T: \Omega \times X \rightarrow CB(X)$ is a random multivalued operator, if for every $x \in X, T(\cdot, x)$ is measurable. A measurable mapping $\xi: \Omega \rightarrow X$ is called random fixed point of a random multivalued operator $T: \Omega \times X \rightarrow CB(X)$ ($f: \Omega \times X \rightarrow X$) if for every $\omega \in \Omega, \xi(\omega) \in T(\omega, \xi(\omega)), f(\omega, \xi(\omega)) =$

$\xi(\omega)$). Let $T: \Omega \times X \rightarrow CB(X)$ be a random operator and $\{\xi_n\}$ a sequence of measurable mappings, $\xi_n: \Omega \rightarrow X$. The sequence $\{\xi_n\}$ is said to be asymptotically T-regular if $d(\xi_n(\omega), T(\omega, \xi_n(\omega))) \rightarrow 0$.

Main Results

Theorem 3.1: Let X be a Polish space. Let $T, S: \Omega \times X \rightarrow CB(X)$ be two continuous random multivalued operators. If there exists measurable mappings $\alpha, \beta: \Omega \rightarrow (0,1)$ such that,

$$\alpha(\omega) H(S(\omega, x), T(\omega, y)) \leq \beta(\omega) \max \begin{pmatrix} d(x, S(\omega, x)), d(y, T(\omega, y)), \\ d(y, S(\omega, x)), d(x, T(\omega, y)), \\ d(x, y) \end{pmatrix} \quad 3.1 (a)$$

For each $x, y \in X, \omega \in \Omega$ and $\alpha, \beta \in R^+$ with $\alpha(\omega) > 1, 0 \leq \beta(\omega) < \alpha(\omega)$, there exists a common random fixed point of S and T . (hence H represents the Hausdroff metric on $CB(X)$ induced by the metric d)

Proof: Let $\xi_0: \Omega \rightarrow X$ be an arbitrary measurable mapping and choose a measurable mapping $\xi_1: \Omega \rightarrow X$ such that $\xi_1(\omega) \in S(\omega, \xi_0(\omega))$ for each $\omega \in \Omega$. then for each $\omega \in \Omega$.

$$\alpha(\omega) H(S(\omega, \xi_0(\omega)), T(\omega, \xi_1(\omega))) \leq \beta(\omega) \max \begin{pmatrix} d(\xi_0(\omega), S(\omega, \xi_0(\omega))), d(\xi_1(\omega), T(\omega, \xi_1(\omega))), \\ d(\xi_1(\omega), S(\omega, \xi_0(\omega))), d(\xi_0(\omega), T(\omega, \xi_1(\omega))), \\ d(\xi_0(\omega), \xi_1(\omega)) \end{pmatrix}$$

Further there exists a measurable mapping $\xi_2: \Omega \rightarrow X$ such that for all $\omega \in \Omega, \xi_2(\omega) \in T(\omega, \xi_1(\omega))$ and

$$\alpha(\omega) d(\xi_1(\omega), \xi_2(\omega)) \leq \beta(\omega) \max \begin{pmatrix} d(\xi_0(\omega), \xi_1(\omega)), d(\xi_1(\omega), \xi_2(\omega)), \\ 0, d(\xi_0(\omega), \xi_1(\omega)) + d(\xi_1(\omega), \xi_2(\omega)) \end{pmatrix}$$

There are four cases:

Case – 1: if we take $\max \begin{pmatrix} d(\xi_0(\omega), \xi_1(\omega)), d(\xi_1(\omega), \xi_2(\omega)), \\ 0, d(\xi_0(\omega), \xi_1(\omega)) + d(\xi_1(\omega), \xi_2(\omega)) \end{pmatrix} = 0$ then

$$\alpha(\omega) d(\xi_1(\omega), \xi_2(\omega)) \leq 0$$

Which is contradiction.

Case – 2: if we take $\max \begin{pmatrix} d(\xi_0(\omega), \xi_1(\omega)), d(\xi_1(\omega), \xi_2(\omega)), \\ 0, d(\xi_0(\omega), \xi_1(\omega)) + d(\xi_1(\omega), \xi_2(\omega)) \end{pmatrix} = d(\xi_0(\omega), \xi_1(\omega))$ then

$$\alpha(\omega) d(\xi_1(\omega), \xi_2(\omega)) \leq \beta(\omega) d(\xi_0(\omega), \xi_1(\omega))$$

Case – 3: if we take $\max \begin{pmatrix} d(\xi_0(\omega), \xi_1(\omega)), d(\xi_1(\omega), \xi_2(\omega)), \\ 0, d(\xi_0(\omega), \xi_1(\omega)) + d(\xi_1(\omega), \xi_2(\omega)) \end{pmatrix} = d(\xi_1(\omega), \xi_2(\omega))$ then

$$\alpha(\omega) d(\xi_1(\omega), \xi_2(\omega)) \leq \beta(\omega) d(\xi_1(\omega), \xi_2(\omega))$$

$$d(\xi_1(\omega), \xi_2(\omega)) \leq \frac{\beta(\omega)}{\alpha(\omega)} d(\xi_1(\omega), \xi_2(\omega)).$$

Which is contradiction

Case – 4: if we take $\max \begin{pmatrix} d(\xi_0(\omega), \xi_1(\omega)), d(\xi_1(\omega), \xi_2(\omega)), \\ 0, d(\xi_0(\omega), \xi_1(\omega)) + d(\xi_1(\omega), \xi_2(\omega)) \end{pmatrix} = d(\xi_0(\omega), \xi_1(\omega)) + d(\xi_1(\omega), \xi_2(\omega))$ then

$$(\alpha(\omega) - 1) d(\xi_1(\omega), \xi_2(\omega)) \leq \beta(\omega) d(\xi_0(\omega), \xi_1(\omega))$$

$$d(\xi_1(\omega), \xi_2(\omega)) \leq \frac{\beta(\omega)}{(\alpha(\omega)-1)} d(\xi_0(\omega), \xi_1(\omega)).$$

From the above four cases, we have, $\max \left\{ \frac{\beta(\omega)}{(\alpha(\omega)-1)}, \frac{\beta(\omega)}{\alpha(\omega)} \right\} = k$ then $d(\xi_1(\omega), \xi_2(\omega)) \leq k d(\xi_0(\omega), \xi_1(\omega))$

By Beg and Shahzad, we obtain a measurable mapping $\xi_3 : \Omega \rightarrow X$ such that for all $\omega \in \Omega$, $\xi_3(\omega) \in S(\omega, \xi_2(\omega))$ and

$$\alpha(\omega) d(\xi_2(\omega), \xi_3(\omega)) \leq \beta(\omega) \max \left\{ \begin{array}{l} d(\xi_1(\omega), \xi_2(\omega)), d(\xi_2(\omega), \xi_3(\omega)), \\ 0, d(\xi_1(\omega), \xi_2(\omega)) + d(\xi_2(\omega), \xi_3(\omega)) \end{array} \right\}$$

There are four cases:

Case – 1: if we take $\max \left\{ \begin{array}{l} d(\xi_1(\omega), \xi_2(\omega)), d(\xi_2(\omega), \xi_3(\omega)), \\ 0, d(\xi_1(\omega), \xi_2(\omega)) + d(\xi_2(\omega), \xi_3(\omega)) \end{array} \right\} = 0$ then

$$\alpha(\omega) d(\xi_2(\omega), \xi_3(\omega)) \leq 0$$

Which is contradiction.

Case – 2: if we take $\max \left\{ \begin{array}{l} d(\xi_1(\omega), \xi_2(\omega)), d(\xi_2(\omega), \xi_3(\omega)), \\ 0, d(\xi_1(\omega), \xi_2(\omega)) + d(\xi_2(\omega), \xi_3(\omega)) \end{array} \right\} = d(\xi_1(\omega), \xi_2(\omega))$ then

$$\alpha(\omega) d(\xi_2(\omega), \xi_3(\omega)) \leq \beta(\omega) d(\xi_1(\omega), \xi_2(\omega))$$

Case – 3: if we take $\max \left\{ \begin{array}{l} d(\xi_1(\omega), \xi_2(\omega)), d(\xi_2(\omega), \xi_3(\omega)), \\ 0, d(\xi_1(\omega), \xi_2(\omega)) + d(\xi_2(\omega), \xi_3(\omega)) \end{array} \right\} = d(\xi_2(\omega), \xi_3(\omega))$ then

$$\alpha(\omega) d(\xi_2(\omega), \xi_3(\omega)) \leq \beta(\omega) d(\xi_2(\omega), \xi_3(\omega))$$

$$d(\xi_2(\omega), \xi_3(\omega)) \leq \frac{\beta(\omega)}{\alpha(\omega)} d(\xi_2(\omega), \xi_3(\omega)).$$

Which is contradiction

Case – 4: if we take $\max \left\{ \begin{array}{l} d(\xi_1(\omega), \xi_2(\omega)), d(\xi_2(\omega), \xi_3(\omega)), \\ 0, d(\xi_1(\omega), \xi_2(\omega)) + d(\xi_2(\omega), \xi_3(\omega)) \end{array} \right\} = d(\xi_1(\omega), \xi_2(\omega)) + d(\xi_2(\omega), \xi_3(\omega))$ then

$$(\alpha(\omega) - 1) d(\xi_2(\omega), \xi_3(\omega)) \leq \beta(\omega) d(\xi_1(\omega), \xi_2(\omega))$$

$$d(\xi_2(\omega), \xi_3(\omega)) \leq \frac{\beta(\omega)}{(\alpha(\omega)-1)} d(\xi_1(\omega), \xi_2(\omega)).$$

From the above four cases, we have, $\max \left\{ \frac{\beta(\omega)}{(\alpha(\omega)-1)}, \frac{\beta(\omega)}{\alpha(\omega)} \right\} = k$ then

$$d(\xi_2(\omega), \xi_3(\omega)) \leq k d(\xi_1(\omega), \xi_2(\omega))$$

Similarly, proceeding the same way, by induction, we get a sequence of measurable mapping $\xi_n : \Omega \rightarrow X$ such that for $n > 0$ and for any $\omega \in \Omega$,

$$\xi_{2n+1}(\omega) \in S(\omega, \xi_{2n}(\omega)), \text{ and } \xi_{2n+2}(\omega) \in T(\omega, \xi_{2n+1}(\omega))$$

This gives, $d(\xi_n(\omega), \xi_{n+1}(\omega)) \leq kd(\xi_{n-1}(\omega), \xi_n(\omega)) \leq \dots \dots \dots \leq k^n d(\xi_0(\omega), \xi_1(\omega))$

For any $m, n \in \mathbb{N}$ such that $m > n$, also by using triangular inequality we have $d(\xi_n(\omega), \xi_m(\omega)) \leq \frac{k^n}{1-k} d(\xi_0(\omega), \xi_1(\omega))$

Which tends to zero as $n \rightarrow \infty$. It follows that $\{\xi_n(\omega)\}$ is a Cauchy sequence and there exists a measurable mapping $\xi : \Omega \rightarrow X$ such that $\xi_n(\omega) \rightarrow \xi(\omega)$ for each $\omega \in \Omega$. It implies that $\xi_{2n+1}(\omega) \rightarrow \xi(\omega)$. Thus we have for any $\omega \in \Omega$,

$$\begin{aligned} d(\xi(\omega), S(\omega, \xi(\omega))) &\leq d(\xi(\omega), \xi_{2n+2}(\omega)) + d(\xi(\omega), S(\omega, \xi_{2n+2}(\omega))) \\ d(\xi(\omega), S(\omega, \xi(\omega))) &\leq d(\xi(\omega), \xi_{2n+2}(\omega)) + H(T(\omega, \xi_{2n+1}(\omega)), S(\omega, \xi_{2n+2}(\omega))) \end{aligned}$$

Therefore, $\alpha(\omega)d(\xi(\omega), S(\omega, \xi(\omega))) \leq d(\xi(\omega), \xi_{2n+2}(\omega)) +$

$$\beta(\omega) \max \left\{ \begin{aligned} &d(\xi_{2n+2}(\omega), S(\omega, \xi_{2n+2}(\omega))), d(\xi_{2n+1}(\omega), T(\omega, \xi_{2n+1}(\omega))), \\ &d(\xi_{2n+1}(\omega), S(\omega, \xi_{2n+2}(\omega))), d(\xi_{2n+2}(\omega), T(\omega, \xi_{2n+1}(\omega))), \\ &d(\xi(\omega), (\omega, \xi_{2n+1}(\omega))) \end{aligned} \right\}$$

Taking as $n \rightarrow \infty$, we have $\alpha(\omega)d(\xi(\omega), S(\omega, \xi(\omega))) \leq \beta(\omega)d(\xi(\omega), S(\omega, \xi(\omega)))$

Which contradiction, hence $\xi(\omega) = S(\omega, \xi(\omega))$ for all $\omega \in \Omega$.

Similarly, for any $\omega \in \Omega$,

$$d(\xi(\omega), S(\omega, \xi(\omega))) \leq d(\xi(\omega), \xi_{2n+1}(\omega)) + H(S(\omega, \xi_{2n}(\omega)), T(\omega, \xi_{2n+1}(\omega)))$$

Hence $\xi(\omega) = T(\omega, \xi(\omega))$ for all $\omega \in \Omega$.

It is easy to see that, $\xi(\omega)$ is common fixed point for S and T in X .

Uniqueness:

Let us assume that, $\xi^*(\omega)$ is another fixed point of S and T in X , different from $\xi(\omega)$, then we have

$$d(\xi(\omega), \xi^*(\omega)) \leq d(\xi(\omega), S(\omega, \xi_{2n}(\omega))) + H(S(\omega, \xi_{2n}(\omega)), T(\omega, \xi_{2n+1}(\omega))) + d(T(\omega, \xi_{2n+1}(\omega)), \xi^*(\omega))$$

By using 3.1(a) and $n \rightarrow \infty$ we have, $d(\xi(\omega), \xi^*(\omega)) \leq 0$

Which contradiction,

So we have, $\xi(\omega)$ is unique common fixed point of S and T in X .

Corollary 3.2:- Let X be a Polish space. Let $S^p, T^q : \Omega \times X \rightarrow CB(X)$ be two continuous random multivalued operators. If there exists measurable mappings $\alpha, \beta : \Omega \rightarrow (0,1)$ such that,

$$\alpha(\omega) H(S(\omega, x), T(\omega, y)) \leq \beta(\omega) \max \left\{ \begin{aligned} &d(x, S(\omega, x)), d(y, T(\omega, y)), \\ &d(y, S(\omega, x)), d(x, T(\omega, y)), \\ &d(x, y) \end{aligned} \right\} \tag{3.2(a)}$$

For each $x, y \in X, \omega \in \Omega$ and $\alpha, \beta \in R^+$ with $\alpha(\omega) > 1, 0 \leq \beta(\omega) < \alpha(\omega)$, there exists a common random fixed point of S and T . (hence H represents the Hausdroff metric on $CB(X)$ induced by the metric d)

Proof: From the theorem 3.1, and on taking $p = q = 1$ it is immediate to see that, the corollary is true. If not then we choose a $\xi_0 : \Omega \rightarrow X$ be an arbitrary measurable mapping and choose a measurable mapping $\xi_1 : \Omega \rightarrow X$ such that $\xi_1(\omega) \in S(\omega, \xi_0(\omega))$ for each $\omega \in \Omega$. then for each $\omega \in \Omega$, and by using 3.2(a) the result is follows.

Now our next result is generalization of our previous theorem 3.1, in fact we prove the following theorem.

Theorem 3.3: Let X be a Polish space. Let $T, S : \Omega \times X \rightarrow CB(X)$ be two continuous random multivalued operators. If there exists measurable mappings $\alpha, \beta : \Omega \rightarrow (0,1)$ such that,

$$\alpha(\omega) H^2(S(\omega, x), T(\omega, y)) \leq \beta(\omega) \min \left\{ \begin{aligned} &\max\{d^2(x, S(\omega, x)), d^2(y, T(\omega, y))\}, \\ &\max\{d^2(y, S(\omega, x)), d^2(x, T(\omega, y))\} \end{aligned} \right\} \tag{3.3(a)}$$

For each $x, y \in X, \omega \in \Omega$ and $\alpha, \beta \in R^+$ with $\alpha(\omega) > 1, 0 \leq \beta(\omega) < \alpha(\omega)$, there exists a common random fixed point of S and T . (hence H represents the Hausdroff metric on $CB(X)$ induced by the metric d)

Proof: Let $\xi_0 : \Omega \rightarrow X$ be an arbitrary measurable mapping and choose a measurable mapping $\xi_1 : \Omega \rightarrow X$ such that $\xi_1(\omega) \in S(\omega, \xi_0(\omega))$ for each $\omega \in \Omega$. then for each $\omega \in \Omega$.

$$\alpha(\omega)H^2 \left(S \left(\omega, \xi_0(\omega) \right), T \left(\omega, \xi_1(\omega) \right) \right) \leq \beta(\omega) \min \left\{ \begin{array}{l} \max \left\{ d^2 \left(\xi_0(\omega), S \left(\omega, \xi_0(\omega) \right) \right), d^2 \left(\xi_1(\omega), T \left(\omega, \xi_1(\omega) \right) \right) \right\}, \\ \max \left\{ d^2 \left(\xi_1(\omega), S \left(\omega, \xi_0(\omega) \right) \right), d^2 \left(\xi_0(\omega), T \left(\omega, \xi_1(\omega) \right) \right) \right\} \end{array} \right\}$$

Further there exists a measurable mapping $\xi_2 : \Omega \rightarrow X$ such that for all $\omega \in \Omega, \xi_2(\omega) \in T \left(\omega, \xi_1(\omega) \right)$ and

$$\alpha(\omega)d^2 \left(\xi_1(\omega), \xi_2(\omega) \right) \leq \beta(\omega) \min \left\{ \begin{array}{l} \max \left\{ d^2 \left(\xi_0(\omega), \xi_1(\omega) \right), d^2 \left(\xi_1(\omega), \xi_2(\omega) \right) \right\}, \\ \max \left\{ d^2 \left(\xi_1(\omega), \xi_1(\omega) \right), d^2 \left(\xi_0(\omega), \xi_2(\omega) \right) \right\} \end{array} \right\}$$

$$d^2 \left(\xi_1(\omega), \xi_2(\omega) \right) \leq (\alpha(\omega) + \beta(\omega))d^2 \left(\xi_0(\omega), \xi_1(\omega) \right)$$

Let $k = (\alpha(\omega) + \beta(\omega))$ This gives $d^2 \left(\xi_1(\omega), \xi_2(\omega) \right) \leq k d^2 \left(\xi_0(\omega), \xi_1(\omega) \right)$

By Beg and Shahzad, we obtain a measurable mapping $\xi_3 : \Omega \rightarrow X$ such that for all $\omega \in \Omega, \xi_3(\omega) \in S \left(\omega, \xi_2(\omega) \right)$ and by using 3.3 (a), we have $d^2 \left(\xi_2(\omega), \xi_3(\omega) \right) \leq k d^2 \left(\xi_1(\omega), \xi_2(\omega) \right) \leq k^2 d^2 \left(\xi_0(\omega), \xi_1(\omega) \right)$

Similarly, proceeding the same way, by induction, we get a sequence of measurable mapping $\xi_n : \Omega \rightarrow X$ such that for $n > 0$ and for any $\omega \in \Omega$,

$$\xi_{2n+1}(\omega) \in S \left(\omega, \xi_{2n}(\omega) \right), \text{ and } \xi_{2n+2}(\omega) \in T \left(\omega, \xi_{2n+1}(\omega) \right)$$

This gives, $d^2 \left(\xi_n(\omega), \xi_{n+1}(\omega) \right) \leq k d^2 \left(\xi_{n-1}(\omega), \xi_n(\omega) \right) \leq \dots \dots \dots \leq k^n d^2 \left(\xi_0(\omega), \xi_1(\omega) \right)$

For any $m, n \in N$ such that $m > n$, also by using triangular inequality we have

$$d^2 \left(\xi_n(\omega), \xi_m(\omega) \right) \leq \frac{k^n}{1-k} d^2 \left(\xi_0(\omega), \xi_1(\omega) \right)$$

Which tends to zero as $n \rightarrow \infty$. It follows that $\{\xi_n(\omega)\}$ is a Cauchy sequence and there exists a measurable mapping $\xi : \Omega \rightarrow X$ such that $\xi_n(\omega) \rightarrow \xi(\omega)$ for each $\omega \in \Omega$. It implies that $\xi_{2n+1}(\omega) \rightarrow \xi(\omega)$. Thus we have for any $\omega \in \Omega$,

$$d^2 \left(\xi(\omega), S(\omega, \xi(\omega)) \right) \leq d^2 \left(\xi(\omega), \xi_{2n+2}(\omega) \right) + d^2 \left(\xi(\omega), S \left(\omega, \xi_{2n+2}(\omega) \right) \right)$$

$$d^2 \left(\xi(\omega), S(\omega, \xi(\omega)) \right) \leq d^2 \left(\xi(\omega), \xi_{2n+2}(\omega) \right) + H^2 \left(T \left(\omega, \xi_{2n+1}(\omega) \right), S \left(\omega, \xi_{2n+2}(\omega) \right) \right)$$

Therefore, by using 3.3(a) we have $d^2 \left(\xi(\omega), S(\omega, \xi(\omega)) \right) \leq (\alpha(\omega) + \beta(\omega)) d^2 \left(\xi(\omega), S(\omega, \xi(\omega)) \right)$

Which contradiction, hence $\xi(\omega) = S(\omega, \xi(\omega))$ for all $\omega \in \Omega$.

Similarly, for any $\omega \in \Omega$,

$$d^2 \left(\xi(\omega), S(\omega, \xi(\omega)) \right) \leq d^2 \left(\xi(\omega), \xi_{2n+1}(\omega) \right) + H^2 \left(S \left(\omega, \xi_{2n}(\omega) \right), T \left(\omega, \xi_{2n+1}(\omega) \right) \right)$$

Hence $\xi(\omega) = T(\omega, \xi(\omega))$ for all $\omega \in \Omega$.

It is easy to see that, $\xi(\omega)$ is common fixed point for S and T in X .

Uniqueness

Let us assume that, $\xi^*(\omega)$ is another fixed point of S and T in X , different from $\xi(\omega)$, then we have

$$d^2 \left(\xi(\omega), \xi^*(\omega) \right) \leq d^2 \left(\xi(\omega), S \left(\omega, \xi_{2n}(\omega) \right) \right) + H^2 \left(S \left(\omega, \xi_{2n}(\omega) \right), T \left(\omega, \xi_{2n+1}(\omega) \right) \right) + d^2 \left(T \left(\omega, \xi_{2n+1}(\omega) \right), \xi^*(\omega) \right)$$

By using 3.3(a) and $n \rightarrow \infty$ we have, $d^2 \left(\xi(\omega), \xi^*(\omega) \right) \leq 0$

Which contradiction,

So we have, $\xi(\omega)$ is unique common fixed point of S and T in X .

Corollary 3.4 Let X be a Polish space. Let $S^p, T^q : \Omega \times X \rightarrow CB(X)$ be two continuous random multivalued operators. If there exists measurable mappings $\alpha, \beta : \Omega \rightarrow (0,1)$ such that,

$$\alpha(\omega)H^2(S(\omega, x), T(\omega, y)) \leq \beta(\omega) \min \left\{ \begin{array}{l} \max\{d^2(x, S(\omega, x)), d^2(y, T(\omega, y))\}, \\ \max\{d^2(y, S(\omega, x)), d^2(x, T(\omega, y))\} \end{array} \right\} \quad 3.4(a)$$

For each $x, y \in X, \omega \in \Omega$ and $\alpha, \beta \in R^+$ with $\alpha(\omega) > 1, 0 \leq \beta(\omega) < \alpha(\omega)$, there exists a common random fixed point of S and T . (hence H represents the Hausdroff metric on $CB(X)$ induced by the metric d)

Proof: - From the theorem 3.3, and on taking $p = q = 1$ it is immediate to see that, the corollary is true. If not then we choose a $\xi_0 : \Omega \rightarrow X$ be an arbitrary measurable mapping and choose a measurable mapping $\xi_1 : \Omega \rightarrow X$ such that $\xi_1(\omega) \in S(\omega, \xi_0(\omega))$ for each $\omega \in \Omega$. then for each $\omega \in \Omega$, and by using 3.3(a) the result is follows.

Conclusion

In this article we prove Some Common Fixed Point Theorems for Random Operator in polish spaces, by using some new type of contractive condition. In fact our main result is more general then other previous known results..

References

1. Spacek A., Zufallige Gleichungen, *Czechoslovak Math. J.*, **5**, 462-466 (1955)
2. Hans O., Reduzierede, *Czech Math, J.*, **7**, 154-158 (1957)
3. Hans O., Random Operator Equations, Proc. 4th Berkeley Symp. Math. Statist. Probability (1960), **II**, 180-202 (1961)
4. Bharucha – Reid A.T., Random Integral Equations, Academic Press, New York, (1972)
5. Itoh. S., *Pacific J. Math.*, **68**, 85-90 (1977)
6. Seghal V.M., and Singh S.P., *Proc. Amer. Math. Soc.* **95**, 91-94 (1985)
7. Papageorgiou N.S., *Proc. Amer. Math. Soc.*, **97**, 507-514 (1986)
8. Rohades B.E., Sessa S. Khan M.S. and Swaleh M., *J. Austral. Math. Soc. (Ser.A)* **43**, 328-346 (1987)
9. Beg I. and Shahzad N., *Nonlinear Anal.* **20**, 835-347 (1993)
10. Beg I. and Azam A., *J. Austral. Math. Soc. Ser. A.*, **53**, 313-326 (1992)
11. Beg I. and Shahzad N., *J. Appl. Math. and Stoch. Analysis* **6**, 95-106 (1993)
12. Heardy G. E. and Rogers T.D., *Canad. Math. Bull.*, **16**, 201-206 (1973)
13. Kanan R., *Bull. Calcutta Math. Soc.*, **60**, 71-76 (1968)
14. Kuratowski K. and Ryll-Nardzewski C., *Bull. Acad. Polo. Sci. Ser. Sci. Math Astronom. Phys.*, **13**, 397-403 (1965)
15. Lin. T.C., *Proc. Amer. Math. Soc.* **103**, 1129-1135 (1988)
16. Wong C.S., *Paci. J. Math.* **48**, 299-312 (1973)
17. Tan. K.K., Xu, H.K., On Fixed Point Theorems of Non-expansive Mappings in Product Spaces, *Proc. Amer. Math. Soc.* **113**, 983-989 (1991)