



Review Paper

## Role of Bio-Fertilizer in Organic Agriculture: A Review

Mishra D.J.<sup>1</sup>, Singh Rajvir<sup>2</sup>, Mishra U.K.<sup>3</sup> and Shahi Sudhir Kumar<sup>4</sup>

<sup>1</sup>Dept. of Soil Science and Ag. Chemistry, Janta College, Bakewar, Etawah, UP, INDIA

<sup>2</sup>Dept. of Agronomy, Janta College, Bakewar, Etawah, UP, INDIA

<sup>3</sup>Dept. of Ag. Botany, Janta College, Bakewar, Etawah, UP, INDIA

<sup>4</sup>Dept. of A.H. and Dairying, Janta College, Bakewar, Etawah, UP, INDIA

Available online at: [www.isca.in](http://www.isca.in)

Received 7<sup>th</sup> November 2012, revised 27<sup>th</sup> August 2012, accepted 29<sup>th</sup> August 2012

### Abstract

*Bio-fertilizers are one of the best modern tools for agriculture. It is a gift of our modern agricultural science. Biofertilizers are applied in the agricultural field as a replacement to our conventional fertilizers. Conventional fertilizers contain compost; household wastes and green manure. Those are not as effective as chemical fertilizers. So, farmers often try to use chemical fertilizers in the field for crop development. But obviously the chemical fertilizers are not environment friendly. They are responsible for water, air and soil pollution and can spread cancer causing agents. Moreover, they may destroy the fertility of the soil in a long run. Scientists have developed Biofertilizers to prevent pollution and to make this world healthy for everybody in a natural way. Bio-fertilizer contains microorganisms which promote the adequate supply of nutrients to the host plants and ensure their proper development of growth and regulation in their physiology. Living microorganisms are used in the preparation of bio-fertilizers. Only those microorganisms are used which have specific functions to enhance plant growth and reproduction. There are different types of microorganisms which are used in the bio-fertilizers. Bio-fertilizer being essential components of Organic farming play vital role in maintaining long term soil fertility and sustainability.*

**Keywords:** Azospirillum, bio-fertilizer, crop growth, sustainability, VAM.

### Introduction

Organic farming has emerged as an important priority area globally in view of the growing demand for safe and healthy food and long term sustainability and concerns on environmental pollution associated with indiscriminate use of agrochemicals. Though the use of chemical inputs in agriculture is inevitable to meet the growing demand for food in world, there are opportunities in selected crops and niche areas where organic production can be encouraged to tap the domestic export market. Bio-fertilizers are being essential component of organic farming are the preparations containing live or latent cells of efficient strains of nitrogen fixing, phosphate solubilizing or cellulolytic micro-organisms used for application to seed, soil or composting areas with the objective of increasing number of such micro-organisms and accelerate those microbial processes which augment the availability of nutrients that can be easily assimilated by plants. Biofertilizers play a very significant role in improving soil fertility by fixing atmospheric nitrogen, both, in association with plant roots and without it, solubilise insoluble soil phosphates and produces plant growth substances in the soil. They are in fact being promoted to harvest the naturally available, biological system of nutrient mobilization (Venkateshwarlu, 2008). The role and importance of bio-fertilizers in sustainable crop production has been reviewed by several authors. But the progress in the field

of BF production technology remained always below satisfaction in Asia because of various constraints.

### Need of Bio-Fertilizers

Indiscriminate use of synthetic fertilizers has led to the pollution and contamination of the soil, has polluted water basins, destroyed micro-organisms and friendly insects, making the crop more prone to diseases and reduced soil fertility.

Demand is much higher than the availability. It is estimated that by 2020, to achieve the targeted production of 321 million tones of food grain, the requirement of nutrient will be 28.8 million tones, while their availability will be only 21.6 million tones being a deficit of about 7.2 million tones<sup>1</sup>.

Depleting feedstock/fossil fuels (energy crisis) and increasing cost of fertilizers. This is becoming unaffordable by small and marginal farmers, depleting soil fertility due to widening gap between nutrient removal and supplies, growing concern about environmental hazards, increasing threat to sustainable agriculture. Besides above facts, the long term use of bio-fertilizers is economical, eco-friendly, more efficient, productive and accessible to marginal and small farmers over chemical fertilizers<sup>2</sup>.

## Potential Characteristic features of some bio-fertilizers

**Nitrogen fixers *Rhizobium*:** belongs to family *Rhizobiaceae*, symbiotic in nature, fix nitrogen 50-100 kg/ ha in association with legumes only. It is useful for pulse legumes like chickpea, red-gram, pea, lentil, black gram, etc., oil-seed legumes like soybean and groundnut and forage legumes like *berseem* and *lucerne*. Successful nodulation of leguminous crops by *Rhizobium* largely depends on the availability of compatible strain for a particular legume. It colonizes the roots of specific legumes to form tumor like growths called root nodules, which acts as factories of ammonia production. *Rhizobium* has ability to fix atmospheric nitrogen in symbiotic association with legumes and certain non-legumes like *Parasponia*. *Rhizobium* population in the soil depends on the presence of legume crops in the field. In absence of legumes, the population decreases. Artificial seed inoculation is often needed to restore the population of effective strains of the *Rhizobium* near the rhizosphere to hasten N-fixation. Each legume requires a specific species of *Rhizobium* to form effective nodules<sup>3</sup>.

***Azospirillum*:** belongs to family *Spirillaceae*, heterotrophic and associative in nature. In addition to their nitrogen fixing ability of about 20-40 kg/ha, they also produce growth regulating substances. Although there are many species under this genus like, *A.amazonense*, *A.halopraeferens*, *A.brasilense*, but, worldwide distribution and benefits of inoculation have been proved mainly with the *A.lipoferum* and *A.brasilense*. The *Azospirillum* form associative symbiosis with many plants particularly with those having the C4-dicarboxylic path way of photosynthesis (Hatch and Slack pathway), because they grow and fix nitrogen on salts of organic acids such as malic, aspartic acid. Thus it is mainly recommended for maize, sugarcane, sorghum, pearl millet etc. The *Azotobacter* colonizing the roots not only remains on the root surface but also a sizable proportion of them penetrates into the root tissues and lives in harmony with the plants. They do not, however, produce any visible nodules or out growth on root tissue.

***Azotobacter*:** belongs to family *Azotobacteriaceae*, aerobic, free living, and heterotrophic in nature. *Azotobacters* are present in neutral or alkaline soils and *A. chroococcum* is the most commonly occurring species in arable soils. *A. vinelandii*, *A. beijerinckii*, *A. insignis* and *A. macrocytogenes* are other reported species. The number of *Azotobacter* rarely exceeds of 104 to 105 g-1 of soil due to lack of organic matter and presence of antagonistic microorganisms in soil. The bacterium produces anti-fungal antibiotics which inhibits the growth of several pathogenic fungi in the root region thereby preventing seedling mortality to a certain extent. The population of *Azotobacter* is generally low in the rhizosphere of the crop plants and in uncultivated soils. The occurrence of this organism has been reported from the rhizosphere of a number of crop plants such as rice, maize, sugarcane, bajra, vegetables and plantation crops.

**Blue Green Algae (*Cyanobacteria*) and *Azolla*:** These belongs to eight different families, phototrophic in nature and produce Auxin, Indole acetic acid and Gibberllic acid, fix 20-30 kg N/ha in submerged rice fields as they are abundant in paddy, so also referred as 'paddy organisms'. N is the key input required in large quantities for low land rice production. Soil N and BNF by associated organisms are major sources of N for low land rice<sup>4</sup>. The 50-60% N requirement is met through the combination of mineralization of soil organic N and BNF by free living and rice plant associated bacteria. To achieve food security through sustainable agriculture, the requirement for fixed nitrogen must be increasingly met by BNF rather than by industrial nitrogen fixation.

BGA forms symbiotic association capable of fixing nitrogen with fungi, liverworts, ferns and flowering plants, but the most common symbiotic association has been found between a free floating aquatic fern, the *Azolla* and *Anabaena azollae* (BGA). *Azolla* contains 4-5% N on dry basis and 0.2-0.4% on wet basis and can be the potential source of organic manure and nitrogen in rice production. The important factor in using *Azolla* as biofertilizer for rice crop is its quick decomposition in the soil and efficient availability of its nitrogen to rice plants. Besides N-fixation, these biofertilizers or biomanures also contribute significant amounts of P, K, S, Zn, Fe, Mb and other micronutrient. The fern forms a green mat over water with a branched stem, deeply bilobed leaves and roots. The dorsal fleshy lobe of the leaf contains the algal symbiont within the central cavity. *Azolla* can be applied as green manure by incorporating in the fields prior to rice planting. The most common species occurring in India is *A. pinnata* and same can be propagated on commercial scale by vegetative means. It may yield on average about 1.5 kg per square meter in a week. India has recently introduced some species of *Azolla* for their large biomass production, which are *A.caroliniana*, *A. microphylla*, *A. filiculoides* and *A. mexicana*.

**Phosphate solubilizers:** Several reports have examined the ability of different bacterial species to solubilize insoluble inorganic phosphate compounds, such as tricalcium phosphate, dicalcium phosphate, hydroxyapatite, and rock phosphate. Among the bacterial genera with this capacity are *pseudomonas*, *Bacillus*, *Rhizobium*, *Burkholderia*, *Achromobacter*, *Agrobacterium*, *Micrococcus*, *Aereobacter*, *Flavobacterium* and *Erwinia*. There are considerable populations of phosphatesolubilizing bacteria in soil and in plant rhizospheres. These include both aerobic and anaerobic strains, with a prevalence of aerobic strains in submerged soils. A considerably higher concentration of phosphate solubilizing bacteria is commonly found in the rhizosphere in comparison with non rhizosphere soil. The soil bacteria belonging to the genera *Pseudomonas* and *Bacillus* and *Fungi* are more common.

**Phosphate absorbers (*Mycorrhiza*):** The term *Mycorrhiza* denotes "fungus roots". It is a symbiotic association between host plants and certain group of fungi at the root system, in

which the fungal partner is benefited by obtaining its carbon requirements from the photosynthates of the host and the host in turn is benefited by obtaining the much needed nutrients especially phosphorus, calcium, copper, zinc etc., which are otherwise inaccessible to it, with the help of the fine absorbing hyphae of the fungus. These fungi are associated with majority of agricultural crops, except with those crops/plants belonging to families of *Chenopodiaceae*, *Amaranthaceae*, *Caryophyllaceae*, *Polygonaceae*, *Brassicaceae*, *Commelinaceae*, *Juncaceae* and *Cyperaceae*.

**Zinc solubilizers:** The nitrogen fixers like *Rhizobium*, *Azospirillum*, *Azotobacter*, BGA and Phosphate solubilizing bacteria like *B. magaterium*, *Pseudomonas striata*, and phosphate mobilizing Mycorrhiza have been widely accepted as bio-fertilizers. However these supply only major nutrients but a host of microorganism that can transform micronutrients are there in soil that can be used as bio-fertilizers to supply micronutrients like zinc, iron, copper etc., The zinc can be solubilized by microorganisms viz., *B. subtilis*, *Thiobacillus thioxidans* and *Saccharomyces sp.* These microorganisms can be used as bio-fertilizers for solubilization of fixed micronutrients like zinc. The results have shown that a *Bacillus sp.* (Zn solubilizing bacteria) can be used as bio-fertilizer for zinc or in soils where native zinc is higher or in conjunction with insoluble cheaper zinc compounds like zinc oxide (ZnO), zinc carbonate (ZnCO<sub>3</sub>) and zinc sulphide (ZnS) instead of costly zinc sulphate.

### Potential role of bio-fertilizers in agriculture

The incorporation of bio-fertilizers (N-fixers) plays major role in improving soil fertility, yield attributing characters and thereby final yield has been reported by many workers. In addition, their application in soil improves soil biota and minimizes the sole use of chemical fertilizers.

Under temperate conditions, inoculation of *Rhizobium* improved number of pods plant<sup>-1</sup>, number of seed pod<sup>-1</sup> and 1000-seed weight (g) and thereby yield over the control. The number of pods plant<sup>-1</sup>, number of seed pod<sup>-1</sup> and 1000-seed weight. In rice under low land conditions, the application of BGA+ *Azospirillum* proved significantly beneficial in improving LAI and all yield attributing aspects.

It is an established fact that the efficiency of phosphate fertilizers is very low (15-20%) due to its fixation in acidic and alkaline soils and unfortunately both soil types are predominating in India accounting more than 34% acidity affected and more than seven million hectares of productive

land salinity/alkaline affected<sup>5</sup>. Therefore, the inoculations with PSB and other useful microbial inoculants in these soils become mandatory to restore and maintain the effective microbial populations for solubilization of chemically fixed phosphorus and availability of other macro and micronutrients to harvest good sustainable yield of various crops.

### Conclusion

Bio-fertilizers being essential components of organic farming play vital role in maintaining long term soil fertility and sustainability by fixing atmospheric di-nitrogen (N=N), mobilizing fixed macro and micro nutrients or convert insoluble P in the soil into forms available to plants, there by increases their efficiency and availability. Currently there is a gap of ten million tones of plant nutrients between removal of crops and supply through chemical fertilizers. In context of both the cost and environmental impact of chemical fertilizers, excessive reliance on the chemical fertilizers is not viable strategy in long run because of the cost, both in domestic resources and foreign exchange, involved in setting up of fertilizer plants and sustaining the production. In this context, organic manures (bio-fertilizers) would be the viable option for farmers to increase productivity per unit area.

### References

1. Arun K.S., Bio-fertilizers for sustainable agriculture. Mechanism of P-solubilization Sixth edition, *Agribios publishers, Jodhpur, India*, 196-197 (2007)
2. Subba Roa N.S., An appraisal of biofertilizers in India. The biotechnology of biofertilizers, (ed.) S.Kannaiyan, *Narosa Pub. House, New* (2001)
3. Venkateshwarlu B. Role of bio-fertilizers in organic farming: Organic farming in rain fed agriculture: Central institute for dry land agriculture, *Hyderabad*, 85-95 (2008)
4. Wani S.P. and Lee K.K., Microorganisms as biological inputs for sustainable agriculture in Organic Agriculture (Thampan, P.K.ed.) Peekay Tree Crops Development Foundation, Cochin, India, 39-76 (1995)
5. Wani, S.P. and Lee K.K., Population dynamics of nitrogen fixing bacteria associated with pearl millet (*P. americanum* L.), In biotechnology of nitrogen fixation in the tropics, University of Pertanian, Malaysia, 21-30 (2002)