



Short Communication

Occurrence of Black Soldier Fly *Hermetia illucens* (Diptera: Stratiomyidae) in Biocompost

Gujarathi Gayatri R. and Pejaver Madhuri K.
B.N. Bandodkar College of Science, Thane, Maharashtra, INDIA

Available online at: www.isca.in

Received 28th December 2012, revised 26th January 2013, accepted 14th February 2013

Abstract

The macro-organisms in the compost feed upon the semi converted organic matter, enhance the composting and convert humus into cured compost. These include several arthropods particularly insects. Among these insects soldier fly *Hermetia illucens* (Diptera: Stratiomyidae) was observed during the study. This is the first time known record of this fly in compost in India. This paper studies the occurrence and description of the fly.

Keywords: Compost, black soldier fly.

Introduction

The compost fauna is well studied across India as well globally. While the references on micro fauna are abundant, the ones regarding macro-fauna in compost are scanty. In India the diversity and role of these various macro-organisms has been studied¹, in which the occurrence of Black soldier fly in the compost was first noted.

The black soldier fly, *Hermetia illucens* (Linnaeus), was first seen in 1930 in Hilo Sugar Company in Hawaiian Islands². Black soldier fly is a tropical fly indigenous to the whole of the Americas, from the southern tip of Argentina to Boston and Seattle. But during World War II, the black soldier flies spread into Europe, India, Asia and even Australia. It is a sleek looking fly that's often confused with a wasp. However, like most flies, the black soldier flies only have two wings (wasps have four) and does not possess a stinger³.

According to Newton *et. al.*⁴, the black soldier fly is often associated with the outdoors and livestock, usually around decaying organic matter such as animal waste or plant material. Since the black soldier fly larvae or BSFL as it is commonly called, consume decaying matter, they have been used to reduce animal manure in commercial swine and poultry facilities in western countries, but in India the practice is not much common. In fact according to Diener *et.al*⁵ valorization of municipal organic waste through larval feeding activity of black soldier fly constitutes a potential benefit, especially low and middle income countries. In India, BSFL larvae were found in poultry house in Punjab in 2007⁶

Black soldier flies though look very bulky, are not known as a disease vector, and possess no danger. Adult soldier flies are a potential mechanical vector of various pathogens. A more likely

negative interaction would be accidental ingestion of black soldier fly larvae by animals or humans⁷.

In natural breeding, black soldier flies lay their eggs in moist organic material while in urbanized areas the black soldier fly lays eggs in dumpsters or compost, which provide similar odors and nutritional needs to naturally occurring organic matter⁸.

Members of the family Stratiomyidae usually range in color from yellow, green, black or blue, with some having a metallic appearance. Many are mimics of other flying insects, such as bees and wasps.

Black soldier fly adults are bristle less flies having a wasp-like appearance and are black or blue in color. They also have two translucent "windows" located on the first abdominal segment and their scutellum is often conspicuously developed. The wing veins of fly are crowded near costa and more strongly pigmented than those behind while vein C does not entirely surround the wing. Adults range from 15 to 20 mm in length⁹ while BSFL observed during study was 1 inch (25mm) in length (figure 1). The adult's antennae are elongated with three segments, and legs have white coloration near the end of each leg (figure 2).

Mating usually occurs two days after adult emergence from the pupal case. The male black soldier fly intercepts a passing female in mid-air and they descend in copula¹⁰. Male soldier flies utilize lekking (gathering of male for competitive mating display) sites, where they await female soldier flies. These sites are defended against other male soldier flies. When a male intrudes upon the territory of a resting male, the resting male seizes the intruder. After a brief descent, the invading male retreats from it⁸.



Figure-1

Figure-2



Figure-3

Methodology

During present study, the Black soldier fly, *Hermetia illucens* was observed hovering on the compost bin after addition of fresh feedstock. The compost contained vegetable kitchen waste including stalks and peels of vegetables, eggshells, raw and processed leftover vegetarian as well as non vegetarian food which was added daily in random quantity as per the generation of waste. Black soldier flies were observed on the upper strata of the bin and their larva (figure 3) was found buried in the middle layers of the compost. BSF larvae were kept in the compost itself instead of isolation, in order to get them regular food supply and to observe life cycle of soldier fly.

Result and Discussion

During the study, the BSF larvae were voraciously eating the debris, while the adult fly hovered upon the upper strata of the compost for egg laying. The blackish gray larvae consumed almost every bit of green feedstock within hours of adding and helped to keep the compost bin clean.

Conclusion

According to Sheppard *et. al*⁹ and Diener *et. al*³ due to this capability of BSF Larvae they are commonly used in household and manure composting in western countries since long time. But at the same time it should be noticed that the rearing of BSF

is easier in India as larvae flourish more in tropical environment than in colder one⁹, hence composting using BSFL should be recommended in India as well.

Acknowledgement

The authors are thankful to Zoology department of B.N. Bandodkar College of Science for allowing composting in the premises of the department.

References

1. Sagade G.A. and Pejaver M.K., Study of diversity of insect fauna from the household biocompost, *Bionano Frontier*, **3(1)**, 67-70 (2009)
2. DuPont M. W. and Larish L.B., Soldier Fly: Livestock Management, Insect Pests. *Cooperative Extension Service, College of tropical agriculture and Human Resources*. (University of Hawaii, at Manoa) LH 107 (2003)
3. Diener S., Zurbrug C., and Tockner K., Conversion of organic material by black soldier fly larvae: establishing optimal feeding rates. *Waste Manag. Res.*, **27 (4)**, 603-610 (2009)
4. Newton L., Sheppard C., Watson D.W., Burtle G. and Dove R., Using the black soldier fly, *Hermetia illucens*, as a value-added tool for the management of swine manure, Waste Management Programs, North Carolina State University http://www.cals.ncsu.edu/waste_mgt/smithfield_projects/phase2report05/cd,web%20files/A2.pdf (2005)
5. Diener S., Solano N., Gutierrez F., Zurbrugg C., and Tockner K., Biological Treatment of Municipal Solid waste using Black Soldier Fly Larvae, *Waste Biomass Valor*, (2), 357-363 (2011)
6. Ashuma, Bal M., Kaur P., Rath S., and Juyal P., First report of *Hermetia illucens* larvae in poultry houses of Punjab, *Journal of Parasitic Diseases*, **31(2)**, 145-146 (2007)
7. Goddard J., *Physician's Guide to Arthropods of Medical Importance*, 4th Edition CRC Press LLC Boca Raton, Florida (2003)
8. Diclaro II J. W. and Kaufman P.E., Publication # EENY 461 <http://edis.ifas.ufl.edu/in830> (2009)
9. Sheppard D.C., Tomberlin J.K., Joyce J.A., Kiser B.C., and Sumner S.M., Rearing methods for the black soldier fly (Diptera: Stratiomyidae), *Journal of Medical Entomology*, **39**, 695-698 (2002)
10. Tomberlin J.K. and Sheppard D.C., Lekking behavior of the black soldier fly (Diptera: Stratiomyidae), *Florida Entomologist*, **84**, 729-730 (2001)